


# Haemochrom Diagnostica

Competence in Haemostasis Diagnostics and Endotoxin Analytics

## BMA Workshop Anticoagulants



Haemochrom  
Diagnostica

Haemochrom Diagnostica  
[www.haemochrom.se](http://www.haemochrom.se)  
info[at]haemochrom.se  
+46 31 706 2070

## Anticoagulants

Principles of different Anticoagulants

Monitoring

## Anticoagulants


Characteristics of an ideal anticoagulant

- » Fast onset of effect
- » Wide therapeutic window
- » No adverse effects (HIT type II, liver toxicity, bleedings, ...)
- » With oral administration: Similarly high bioavailability
- » Predictable pharmacokinetics
- » Linear dose-effect relationship
- » Easy dose regimen
- » No interactions with other drugs or foods
- » Fast loss of effect after treatment termination
- » Possibility to monitor the anti-coagulatory effect
- » Available and safe antidote

## Anticoagulants

### Vitamin K-Antagonists


- » Synthesis of vitamin K dependent factors



## Anticoagulants


Inhibition by

- AT
- TFPI
- Protein C Pathway





## Anticoagulants

Monitoring: Principle of measurement





## Anticoagulants

Measurement of inhibitors e.g. Antithrombin, Antiplasmin, Anti FXa, ...



## Anticoagulants

Effect of Rivaroxaban and Dabigatran on aPTT



Andreas Hillarp, Malmö, Equalis Meeting 2009